




Contents lists available at ScienceDirect

Bone

journal homepage: www.elsevier.com/locate/bone

Correspondence

On epidemiology of fractures and variation with age and ethnicity

Keywords:
Osteoporosis
Epidemiology
Fracture
Incidence
CRPD
Ethnicity

We recently demonstrated differences in fracture incidence by ethnicity in the UK population using the Clinical Practice Research Datalink (CPRD) [1]. In their letter, Harper et al. [2] draw attention to the differential age structure of the UK population by ethnicity, and suggest that the lower incidence of fractures in Asian adults over 50 years might reflect the lower proportion of very elderly individuals within this demographic compared with the white population. The relatively small number of fractures within black and South Asian ethnicities meant that presentation of fracture incidence stratified by age band was not feasible. We have thus used a Poisson model to calculate incidence rate ratios (with white ethnicity as the referent), adjusting for age and calendar year to account for any contribution of age distribution and secular change within ethnicity strata. These data are summarised in Table 1 below, and clearly demonstrate that the differences in fracture incidence by ethnicity are congruent with the raw incidence data presented in our paper, with rates highest in whites, intermediate in South Asians and lowest in blacks. In conclusion, whilst there are ethnic

differences in age distribution in the UK, these do not explain the differential fracture rates; as discussed in our paper [1], variations in factors such as BMD, bone geometry and body composition may all contribute. Whatever the underlying cause, we agree with Harper et al. that our findings inform the allocation of healthcare resources to those at greatest need.

References

- [1] E.M. Curtis, R. van der Velde, R.J. Moon, J.P. van den Bergh, P. Geusens, F. de Vries, T.P. van Staa, C. Cooper, N.C. Harvey, Epidemiology of fractures in the United Kingdom 1988-2012: Variation with age, sex, geography, ethnicity and socioeconomic status, *Bone* 87 (2016) 19-26.
- [2] B. Bogin, On epidemiology of fractures and variation with age and ethnicity, 2016.

Nicholas C. Harvey
MRC Lifecourse Epidemiology Unit, University of Southampton,
Southampton SO16 6YD, UK
NIHR Southampton Biomedical Research Centre, University of
Southampton and University Hospital Southampton NHS Foundation Trust,
Tremona Road, Southampton SO16 6YD, UK

Elizabeth M. Curtis
MRC Lifecourse Epidemiology Unit, University of Southampton,
Southampton SO16 6YD, UK

Robert van der Velde
Dept. of Internal Medicine, VieCuri Medical Center, Venloseweg 595971 PB
Venlo, The Netherlands

Rebecca J. Moon
MRC Lifecourse Epidemiology Unit, University of Southampton,
Southampton SO16 6YD, UK
Paediatric Endocrinology, Southampton University Hospitals NHS
Foundation Trust, Southampton SO16 6YD, UK

Joop P.W. van den Bergh
Dept. of Internal Medicine, VieCuri Medical Center, Venloseweg 595971 PB
Venlo, The Netherlands
Maastricht University Medical Center, Maastricht, The Netherlands

Piet Geusens
Department of Internal Medicine, Subdivision of Rheumatology, Maastricht
University Medical Centre, Maastricht, The Netherlands
University Hasselt, Hasselt, Belgium

Frank de Vries
Department of Clinical Pharmacy & Toxicology, Maastricht University
Medical Centre, Maastricht, The Netherlands

Tjeerd P. van Staa
Health eResearch Centre, Farr Institute for Health Informatics Research,
University of Manchester, 1.003 Vaughan House, Portsmouth, Road, M13
9PL, UK

Table 1

Incidence of fragility fracture by ethnicity [1], and Incidence Rate Ratios relative to white ethnicity (adjusted for age and calendar year) in UK men and women aged 50 years or over, 1988 to 2012.

Women			
Ethnicity	Number fractures	Rate (/10,000py)	IRR*
White	71,081	104.03	Ref
Other	496	70.98	0.81 (0.74, 0.89)
Mixed	77	50.12	0.63 (0.51, 0.79)
South Asian	491	44.46	0.57 (0.52, 0.62)
Black	149	22.26	0.28 (0.24, 0.33)
Men			
Ethnicity	Number fractures	Rate (/10,000py)	IRR*
White	25,285	41.74	Ref
Other	170	28.88	0.76 (0.66, 0.89)
South Asian	289	26.85	0.73 (0.65, 0.82)
Mixed	22	17.37	0.49 (0.32, 0.74)
Black	88	15.75	0.42 (0.34, 0.51)

* IRR = Incidence Rate Ratio from Poisson model, adjusting for age and calendar year.

Department of Pharmacoepidemiology & Clinical Pharmacology, University of Utrecht, Utrecht, The Netherlands, 3508, TB

Cyrus Cooper

MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton SO16 6YD, UK

NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Tremona Road, Southampton SO16 6YD, UK

NIHR Musculoskeletal Biomedical Research Unit, University of Oxford, Nuffield Orthopedic Centre, Headington, Oxford OX3 7HE, UK

Corresponding author at: MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton SO16 6YD, UK.

E-mail address: cc@mrc.soton.ac.uk.

14 July 2016

Available online xxxx